

Soil Analysis

Submitted By: **BN88888**
AgSource Test Account
106 NORTH CECIL STREET
PO BOX 7
BONDUEL, WI 54107

Submitted For:
AgSource UW Recs Report Example



Laboratory Sample #
BH69403 - BH69415
 Information Sheet #
S0807-018

Date Received:
08/02/2018

Date Processed:
08/03/2018

Farm Id: Dads Water Str

County: Account No:
 Marathon BN88888
Field: Test Field 1
Acres 40.1
 Soil Name/Subsoil group:
 unknown
 Plow Depth: Previous Crop:
 7.00
 Slope: Irrigated: Tiled:
 No No No

NUTRIENT RECOMMENDATIONS												
Cropping Sequence	Yield Goal	Crop Nutrient Need			Fertilizer Credits				Nutrients to Apply			
		N	P ₂ O ₅	K ₂ O	Legume N	Manure N	P ₂ O ₅	K ₂ O	N	P ₂ O ₅	K ₂ O	
	- per acre -	----- lbs/a -----			--- lbs/a ---	----- lbs/a -----				----- lbs/a -----		
Corn, grain	191-210 bu	***	0	15	0	0	0	0	***	0	15	
Corn, grain	171-190 bu	***	0	15	0	0	0	0	***	0	15	
Soybean, grain	76-85 bu	0	0	30	0	0	0	0	0	0	30	
Wheat, grain + straw	81-100 bu	0	0	25	0	0	0	0	0	0	25	

There is no lime recommendation for this rotation. Please see Additional Information below.

*** Please use the new Wisconsin Nitrogen Application Rates table to determine the N Application rate. Table included at end of report.

TEST INTERPRETATION						
Cropping Sequence	Very Low	Low	Optimum	High	Very High	Excessive
P	[Red bar]					
K	[Yellow bar]					
Rotation pH	[Red bar]					

LABORATORY ANALYSIS											LAB USE			MISC							
Adjusted Avg: 7.5 3.6 47 182 1949 515											14.4	3.5	67.0	29.6	100.0						
Sample ID	Soil pH	O.M. %	Phosphorus PPM	Potassium PPM	60-69 Lime Req T/a	Calcium PPM	Magnesium PPM	Boron PPM	Manganese PPM	Zinc PPM	Sulfate Sulfur	Sulfur Avail Index	Texture Code	Sample Density	Buffer Code	Total CEC	% Base Saturation				
																	%K	%Ca	%Mg	Tot %	%H
Test 22	7.5	2.9	42	186		1538	427						2	0.98		11.7	4.1	65.9	30.0	100.0	
Test 23	7.6	2.7	53	221		1596	449						2	0.91		12.2	4.6	65.3	30.1	100.0	
Test 24	7.7	3.4	58	164		1917	501						2	0.85		14.1	3.0	67.9	29.1	100.0	
Test 25	7.4	3.2	80	230		1529	400						2	0.91		11.5	5.1	66.4	28.5	100.0	
Test 17	7.4	3.1	40	157		1770	496						2	0.86		13.3	3.0	66.5	30.5	100.0	
Test 18	7.7	7.6	31	114		3472	705						2	0.76		23.4	1.2	74.1	24.7	100.0	
Test 19	7.7	2.9	22	157		2115	623						2	1.01		16.1	2.5	65.8	31.8	100.0	
Test 20	7.1	3.7	52	240		1908	548						2	0.92		14.6	4.2	65.1	30.7	100.0	
Test 21	7.6	2.8	42	172		1693	485						2	0.94		12.9	3.4	65.7	30.9	100.0	

SECONDARY & MICRONUTRIENT RECOMMENDATIONS

Interpretations -----> Ca-H Mg-H
 Response to added Ca is unlikely.
 Response to added Mg is unlikely.

ADDITIONAL INFORMATION

N.R.=Not required for calculation of lime requirement when soil pH is 6.6 or higher.
 Starter fertilizer (e.g. 10+20+20 lbs N+P₂O₅+K₂O/a) is advisable for row crops on soils slow to warm in the spring.
 Recommended rates are the total amount of nutrients to apply (N-P-K), including starter fertilizer.
 Year 1,2 If corn is harvested for silage instead of grain apply extra 90 lbs K₂O per acre to next crop.
 A lime recommendation is calculated only when soil pH is more than 0.2 units below the optimum pH. Starter fertilizer (e.g. 10 + 20 + 20 lbs N + P₂O₅ + K₂O/a) is advisable for row crops on soils slow to warm in the spring.
 A soil nitrate test may better estimate actual corn N needs. If conservative tillage leaves more than 50% residue cover when corn follows after corn, add an additional 30 N lb/a.
 If alfalfa will be maintained for more than three years, increase recommended: K₂O by 20% each year.

Soil Analysis

Submitted By: **BN88888**
AgSource Test Account
106 NORTH CECIL STREET
PO BOX 7
BONDUEL, WI 54107

Submitted For:
AgSource UW Recs Report Example



Laboratory Sample #
BH69403 - BH69415
 Information Sheet #
S0807-018

Date Received:
08/02/2018

Date Processed:
08/03/2018

Farm Id: Dads Water Str

County: Account No:
 Marathon BN88888
Field: Test Field 10
Acres 9.3
 Soil Name/Subsoil group:
 unknown
 Plow Depth: Previous Crop:
 7.00
 Slope: Irrigated: Tiled:
 No No No

NUTRIENT RECOMMENDATIONS											
Cropping Sequence	Yield Goal	Crop Nutrient Need			Fertilizer Credits				Nutrients to Apply		
		N	P ₂ O ₅	K ₂ O	Legume N	Manure N	P ₂ O ₅	K ₂ O	N	P ₂ O ₅	K ₂ O
	- per acre -	----- lbs/a -----			--- lbs/a ---	----- lbs/a -----			----- lbs/a -----		
Corn, grain	191-210 bu	***	0	15	0	0	0	0	***	0	15
Corn, grain	171-190 bu	***	0	15	0	0	0	0	***	0	15
Soybean, grain	76-85 bu	0	0	30	0	0	0	0	0	0	30
Wheat, grain + straw	81-100 bu	0	0	25	0	0	0	0	0	0	25

There is no lime recommendation for this rotation. Please see Additional Information below.

*** Please use the new Wisconsin Nitrogen Application Rates table to determine the N Application rate. Table included at end of report.

TEST INTERPRETATION						
Cropping Sequence	Very Low	Low	Optimum	High	Very High	Excessive
P	[Bar chart showing P level in the 'Very Low' to 'Low' range]					
K	[Bar chart showing K level in the 'Very High' to 'Excessive' range]					
Rotation pH	[Bar chart showing pH level in the 'Optimum' to 'High' range]					

LABORATORY ANALYSIS											LAB USE				MISC						
Adjusted Avg: 7.6 4.3 46 211 2246 600															16.7 3.2 67.3 29.5 100.0						
Sample ID	Soil pH	O.M. %	Phosphorus PPM	Potassium PPM	60-69 Lime Req T/a	Calcium PPM	Magnesium PPM	Boron PPM	Manganese PPM	Zinc PPM	Sulfate Sulfur	Sulfur Avail Index	Texture Code	Sample Density	Buffer Code	Total CEC	% Base Saturation				
																	%K	%Ca	%Mg	Tot %	%H
Test 15	7.5	4.6	43	195		2270	575						2	0.86		16.6	3.0	68.5	28.5	100.0	
Test 16	7.7	4.0	49	226		2222	625						2	0.92		16.8	3.4	66.1	30.5	100.0	

SECONDARY & MICRONUTRIENT RECOMMENDATIONS										
Interpretations -----> Ca-H Mg-H										
Response to added Ca is unlikely.										
Response to added Mg is unlikely.										

ADDITIONAL INFORMATION

N.R.=Not required for calculation of lime requirement when soil pH is 6.6 or higher.
 Starter fertilizer (e.g. 10+20+20 lbs N+P₂O₅+K₂O/a) is advisable for row crops on soils slow to warm in the spring.
 Recommended rates are the total amount of nutrients to apply (N-P-K), including starter fertilizer.
 Year 1,2 If corn is harvested for silage instead of grain apply extra 90 lbs K₂O per acre to next crop.
 A lime recommendation is calculated only when soil pH is more than 0.2 units below the optimum pH. Starter fertilizer (e.g. 10 + 20 + 20 lbs N + P₂O₅ + K₂O/a) is advisable for row crops on soils slow to warm in the spring.
 A soil nitrate test may better estimate actual corn N needs. If conservative tillage leaves more than 50% residue cover when corn follows after corn, add an additional 30 N lb/a.
 If alfalfa will be maintained for more than three years, increase recommended: K₂O by 20% each year.

Soil Analysis

Submitted By: **BN88888**
AgSource Test Account
106 NORTH CECIL STREET
PO BOX 7
BONDUEL, WI 54107

Submitted For:
AgSource UW Recs Report Example



Laboratory Sample #
BH69403 - BH69415
 Information Sheet #
S0807-018

Date Received:
08/02/2018

Date Processed:
08/03/2018

Farm Id: Dads Water Str

County: Account No:
 Marathon BN88888
Field: Test Field 12
Acres 6.4
 Soil Name/Subsoil group:
 unknown
 Plow Depth: Previous Crop:
 7.00
 Slope: Irrigated: Tiled:
 No No No

NUTRIENT RECOMMENDATIONS											
Cropping Sequence	Yield Goal	Crop Nutrient Need			Fertilizer Credits				Nutrients to Apply		
		N	P ₂ O ₅	K ₂ O	Legume N	Manure N	P ₂ O ₅	K ₂ O	N	P ₂ O ₅	K ₂ O
	- per acre -	----- lbs/a -----			--- lbs/a ---	----- lbs/a -----			----- lbs/a -----		
Corn, grain	191-210 bu	***	0	15	0	0	0	0	***	0	15
Corn, grain	171-190 bu	***	0	15	0	0	0	0	***	0	15
Soybean, grain	76-85 bu	0	0	30	0	0	0	0	0	0	30
Wheat, grain + straw	81-100 bu	0	0	25	0	0	0	0	0	0	25

There is no lime recommendation for this rotation. Please see Additional Information below.

*** Please use the new Wisconsin Nitrogen Application Rates table to determine the N Application rate. Table included at end of report.

TEST INTERPRETATION						
Cropping Sequence	Very Low	Low	Optimum	High	Very High	Excessive
P	[Red bar]					
K	[Yellow bar]					
Rotation pH	[Green bar]					

LABORATORY ANALYSIS										LAB USE				MISC							
Adjusted Avg:		7.7	3.9	43	201	2291		639						17.2	3.0	66.6	30.5	100.0			
Sample ID	Soil pH	O.M. %	Phosphorus PPM	Potassium PPM	60-69 Lime Req T/a	Calcium PPM	Magnesium PPM	Boron PPM	Manganese PPM	Zinc PPM	Sulfate Sulfur	Sulfur Avail Index	Texture Code	Sample Density	Buffer Code	Total CEC	% Base Saturation				
																	%K	%Ca	%Mg	Tot %	%H
Test 13	7.6	3.7	39	201		2187	636						2	0.86		16.7	3.1	65.6	31.3	100.0	
Test 14	7.7	4.0	47	200		2395	642						2	0.85		17.7	2.9	67.5	29.7	100.0	

SECONDARY & MICRONUTRIENT RECOMMENDATIONS										
Interpretations -----> Ca-H Mg-H										
Response to added Ca is unlikely.										
Response to added Mg is unlikely.										

ADDITIONAL INFORMATION

N.R.=Not required for calculation of lime requirement when soil pH is 6.6 or higher.
 Starter fertilizer (e.g. 10+20+20 lbs N+P₂O₅+K₂O/a) is advisable for row crops on soils slow to warm in the spring.
 Recommended rates are the total amount of nutrients to apply (N-P-K), including starter fertilizer.
 Year 1,2 If corn is harvested for silage instead of grain apply extra 90 lbs K₂O per acre to next crop.
 A lime recommendation is calculated only when soil pH is more than 0.2 units below the optimum pH. Starter fertilizer (e.g. 10 + 20 + 20 lbs N + P₂O₅ + K₂O/a) is advisable for row crops on soils slow to warm in the spring.
 A soil nitrate test may better estimate actual corn N needs. If conservative tillage leaves more than 50% residue cover when corn follows after corn, add an additional 30 N lb/a.
 If alfalfa will be maintained for more than three years, increase recommended: K₂O by 20% each year.

Nitrogen Application Rate Guidelines for Corn

(For more info, see <http://www.soils.wisc.edu/extension/pubs/A2809.pdf>)

Justification: While the yield response of corn to applied N has not changed, the economics of corn production have. Recently soil fertility specialists in Wisconsin, Minnesota, Iowa, and Illinois have agreed to use the same philosophy to develop N rate guidelines for corn (grain). The philosophy used is based on maximizing return to N fertilizer. The new N rate guidelines were developed as a means to provide growers guidance on how much they might adjust their N application rates and maintain or enhance profitability depending upon their individual farm situation. Research data collected in Wisconsin from research farms and grower fields over a period of 20 years was used to develop the guidelines.

SUGGESTED N APPLICATION RATES FOR CORN(GRAIN) AT DIFFERENT N: CORN PRICE RATIOS

Soil and Previous Crop	N: Corn Price Ratio (\$/lb N:\$/bu)							
	0.05 Rate *3 Range *4		0.10 Rate *3 Range *4		0.15 Rate *3 Range *4		0.20 Rate *3 Range *4	
HIGH YIELD POTENTIAL SOILS Corn, Forage Legumes, Leguminous vegetables, Green manures *5 Soybean, Small grains *6	lb N/a (Total to Apply) *2							
	190	170-210	165	155-180	150	140-160	135	125-150
	140	125-160	120	105-135	105	95-115	95	80-105
MEDIUM YIELD POTENTIAL SOILS Corn, Forage Legumes, Leguminous vegetables, Green manures *5 Soybean, Small grains *6	145	130-160	125	115-140	115	105-125	105	95-110
	130	110-150	100	85-120	85	70-95	70	60-80
IRRIGATED SANDS AND LOAMY SANDS All Crops *5	215	200-230	200	185-210	185	175-195	175	165-185
NON-IRRIGATED SANDS AND LOAMY SANDS All Crops *5	140	130-150	130	120-140	120	110-130	110	100-120

*1 To determine soil yield potential, consult UWEX publication A2809 or contact your county agent or agronomist.

*2 Includes N in starter.

*3 Maximum return to N (MRTN) rate.

*4 Profitability range within \$1/a or MRTN rate.

*5 Subtract N credit for forage legumes, legume vegetables, animal manures, green manures.

*6 Subtract credits for animal manures and second year forage legumes.

Guidelines for choosing an appropriate N application rate for corn (grain)

- 1) If there is more than 50% residue cover at planting, use the upper end of the range.
- 2) For small grains grown on medium and fine textured soils, the mid to low end of the profitable range is the most appropriate.
- 3) If 100% of the N will come from organic sources, use the top end of the range. In addition, up to 20 lb N/a in starter fertilizer may be applied.
- 4) For medium and fine textured soils with: < 2% organic matter, use the high end of the range; > 10% organic matter, use the low end of the range.
- 5) For coarse textured soils with: < 2% organic matter, use the high end of the range; > 2% organic matter, use the mid to low end of the range.
- 6) If there is a likelihood of residual N, then use the low end of the range or use the high end of the range and subtract preplant nitrate test (PPNT) credits.
- 7) For corn following small grains on medium and fine textured soils, the middle to low end of the range is most appropriate.

Nitrogen Application Rate Guidelines for Wheat

(For more info, see <http://www.soils.wisc.edu/extension/pubs/A2809.pdf>)

SUGGESTED N APPLICATION RATES FOR WHEAT AT DIFFERENT N: WHEAT PRICE RATIOS

Loamy Soil and Previous Crop	N: Wheat Price Ratio (\$/lb N:\$/bu)							
	0.05 Rate *3 Range		0.075 Rate *3 Range		0.10 Rate *3 Range		0.125 Rate *3 Range	
	lb N/a (Total to Apply) *1							
Corn *2 : < 50 or no PPNT	75	65-85	70	55-80	60	50-70	55	40-65
Corn : 51 to 100	45	35-55	40	30-50	35	25-40	30	20-35
Corn : > 100	0	0-0	0	0-0	0	0-0	0	0-0
Soybean, Small grains : All *3	55	45-65	50	40-60	45	35-50	40	35-45

*1 On loamy soils with < 2% organic matter, add 30 lb N/a to all rates. On soils with more than 10% organic matter, reduce rates by 30 lb N/a.

Reduce N rates by 10 lb N/a for spring wheat on all soils. No N is required on organic soils. Manure N credits must be subtracted from these values.

*2 If wheat follows a forage legume or leguminous vegetable, use the MRTN rate for wheat following corn with PPNT < 50 and take the legume credit.

*3 Previous crop soybean or small grain: If a PPNT is taken and the PPNT is < 50 lb N/a, use the top end of the profitable range; if the PPNT is 51 to 100 lb N/a, use the bottom end of the profitable range; if the PPNT is > 100 lb N/a, no additional N is needed. Do not take a soybean legume credit.